# 2. Caratheodory's Extension

In the following,  $\Omega$  is a set. Whenever a union of sets is denoted  $\forall$  as opposed to  $\cup$ , it indicates that the sets involved are pairwise disjoint.

**Definition 6** A semi-ring on  $\Omega$  is a subset S of the power set  $P(\Omega)$  with the following properties:

- (i)  $\emptyset \in \mathcal{S}$
- (ii)  $A, B \in \mathcal{S} \Rightarrow A \cap B \in \mathcal{S}$

(iii) 
$$A, B \in \mathcal{S} \Rightarrow \exists n \geq 0, \exists A_i \in \mathcal{S} : A \setminus B = \biguplus_{i=1}^n A_i$$

The last property (iii) says that whenever  $A, B \in \mathcal{S}$ , there is  $n \geq 0$  and  $A_1, \ldots, A_n$  in  $\mathcal{S}$  which are pairwise disjoint, such that  $A \setminus B = A_1 \uplus \ldots \uplus A_n$ . If n = 0, it is understood that the corresponding union is equal to  $\emptyset$ , (in which case  $A \subseteq B$ ).

**Definition 7** A ring on  $\Omega$  is a subset  $\mathcal{R}$  of the power set  $\mathcal{P}(\Omega)$  with the following properties:

- (i)  $\emptyset \in \mathcal{R}$
- (ii)  $A, B \in \mathcal{R} \Rightarrow A \cup B \in \mathcal{R}$
- (iii)  $A, B \in \mathcal{R} \Rightarrow A \setminus B \in \mathcal{R}$

EXERCISE 1. Show that  $A \cap B = A \setminus (A \setminus B)$  and therefore that a ring is closed under pairwise intersection.

EXERCISE 2.Show that a ring on  $\Omega$  is also a semi-ring on  $\Omega$ .

EXERCISE 3.Suppose that a set  $\Omega$  can be decomposed as  $\Omega = A_1 \uplus A_2 \uplus A_3$  where  $A_1, A_2$  and  $A_3$  are distinct from  $\emptyset$  and  $\Omega$ . Define  $\mathcal{S}_1 \stackrel{\triangle}{=} \{\emptyset, A_1, A_2, A_3, \Omega\}$  and  $\mathcal{S}_2 \stackrel{\triangle}{=} \{\emptyset, A_1, A_2 \uplus A_3, \Omega\}$ . Show that  $\mathcal{S}_1$  and  $\mathcal{S}_2$  are semi-rings on  $\Omega$ , but that  $\mathcal{S}_1 \cap \mathcal{S}_2$  fails to be a semi-ring on  $\Omega$ .

EXERCISE 4. Let  $(\mathcal{R}_i)_{i \in I}$  be an arbitrary family of rings on  $\Omega$ , with  $I \neq \emptyset$ . Show that  $\mathcal{R} \stackrel{\triangle}{=} \cap_{i \in I} \mathcal{R}_i$  is also a ring on  $\Omega$ .

EXERCISE 5. Let  $\mathcal{A}$  be a subset of the power set  $\mathcal{P}(\Omega)$ . Define:

$$R(\mathcal{A}) \stackrel{\triangle}{=} \{ \mathcal{R} \text{ ring on } \Omega : \ \mathcal{A} \subseteq \mathcal{R} \}$$

Show that  $\mathcal{P}(\Omega)$  is a ring on  $\Omega$ , and that  $R(\mathcal{A})$  is not empty. Define:

$$\mathcal{R}(\mathcal{A}) \stackrel{\triangle}{=} \bigcap_{\mathcal{R} \in R(\mathcal{A})} \mathcal{R}$$

Show that  $\mathcal{R}(\mathcal{A})$  is a ring on  $\Omega$  such that  $\mathcal{A} \subseteq \mathcal{R}(\mathcal{A})$ , and that it is the smallest ring on  $\Omega$  with such property, (i.e. if  $\mathcal{R}$  is a ring on  $\Omega$  and  $\mathcal{A} \subseteq \mathcal{R}$  then  $\mathcal{R}(\mathcal{A}) \subseteq \mathcal{R}$ ).

**Definition 8** Let  $A \subseteq \mathcal{P}(\Omega)$ . We call **ring generated** by A, the ring on  $\Omega$ , denoted  $\mathcal{R}(A)$ , equal to the intersection of all rings on  $\Omega$ , which contain A.

EXERCISE 6.Let S be a semi-ring on  $\Omega$ . Define the set R of all finite unions of pairwise disjoint elements of S, i.e.

$$\mathcal{R} \stackrel{\triangle}{=} \{A : A = \bigoplus_{i=1}^{n} A_i \text{ for some } n \geq 0, A_i \in \mathcal{S}\}$$

(where if n=0, the corresponding union is empty, i.e.  $\emptyset \in \mathcal{R}$ ). Let  $A= \bigoplus_{i=1}^n A_i$  and  $B= \bigoplus_{j=1}^p B_j \in \mathcal{R}$ :

- 1. Show that  $A \cap B = \bigcup_{i,j} (A_i \cap B_j)$  and that  $\mathcal{R}$  is closed under pairwise intersection.
- 2. Show that if  $p \ge 1$  then  $A \setminus B = \bigcap_{i=1}^p (\biguplus_{i=1}^n (A_i \setminus B_i))$ .
- 3. Show that  $\mathcal{R}$  is closed under pairwise difference.
- 4. Show that  $A \cup B = (A \setminus B) \oplus B$  and conclude that  $\mathcal{R}$  is a ring on  $\Omega$ .
- 5. Show that  $\mathcal{R}(\mathcal{S}) = \mathcal{R}$ .

EXERCISE 7. Everything being as before, define:

$$\mathcal{R}' \stackrel{\triangle}{=} \{A : A = \bigcup_{i=1}^n A_i \text{ for some } n \geq 0, A_i \in \mathcal{S}\}$$

(We do not require the sets involved in the union to be pairwise disjoint). Using the fact that  $\mathcal{R}$  is closed under finite union, show that  $\mathcal{R}' \subseteq \mathcal{R}$ , and conclude that  $\mathcal{R}' = \mathcal{R} = \mathcal{R}(\mathcal{S})$ .

**Definition 9** Let  $A \subseteq \mathcal{P}(\Omega)$  with  $\emptyset \in A$ . We call **measure** on A, any map  $\mu : A \to [0, +\infty]$  with the following properties:

$$(i) \qquad \mu(\emptyset) = 0$$

(ii) 
$$A \in \mathcal{A}, A_n \in \mathcal{A} \text{ and } A = \bigcup_{n=1}^{+\infty} A_n \Rightarrow \mu(A) = \sum_{n=1}^{+\infty} \mu(A_n)$$

The  $\forall$  indicates that we assume the  $A_n$ 's to be pairwise disjoint in the l.h.s. of (ii). It is customary to say in view of condition (ii) that a measure is *countably* additive.

EXERCISE 8.If  $\mathcal{A}$  is a  $\sigma$ -algebra on  $\Omega$  explain why property (ii) can be replaced by:

$$(ii)'$$
  $A_n \in \mathcal{A}$  and  $A = \bigcup_{n=1}^{+\infty} A_n \Rightarrow \mu(A) = \sum_{n=1}^{+\infty} \mu(A_n)$ 

EXERCISE 9. Let  $\mathcal{A} \subseteq \mathcal{P}(\Omega)$  with  $\emptyset \in \mathcal{A}$  and  $\mu : \mathcal{A} \to [0, +\infty]$  be a measure on  $\mathcal{A}$ .

- 1. Show that if  $A_1, \ldots, A_n \in \mathcal{A}$  are pairwise disjoint and the union  $A = \bigcup_{i=1}^n A_i$  lies in  $\mathcal{A}$ , then  $\mu(A) = \mu(A_1) + \ldots + \mu(A_n)$ .
- 2. Show that if  $A, B \in \mathcal{A}, A \subseteq B$  and  $B \setminus A \in \mathcal{A}$  then  $\mu(A) \leq \mu(B)$ .

EXERCISE 10. Let S be a semi-ring on  $\Omega$ , and  $\mu: S \to [0, +\infty]$  be a measure on S. Suppose that there exists an extension of  $\mu$  on  $\mathcal{R}(S)$ , i.e. a measure  $\bar{\mu}: \mathcal{R}(S) \to [0, +\infty]$  such that  $\bar{\mu}_{|S} = \mu$ .

- 1. Let A be an element of  $\mathcal{R}(\mathcal{S})$  with representation  $A = \bigoplus_{i=1}^{n} A_i$  as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . Show that  $\bar{\mu}(A) = \sum_{i=1}^{n} \mu(A_i)$
- 2. Show that if  $\bar{\mu}': \mathcal{R}(\mathcal{S}) \to [0, +\infty]$  is another measure with  $\bar{\mu}'_{|\mathcal{S}} = \mu$ , i.e. another extension of  $\mu$  on  $\mathcal{R}(\mathcal{S})$ , then  $\bar{\mu}' = \bar{\mu}$ .

EXERCISE 11. Let S be a semi-ring on  $\Omega$  and  $\mu: S \to [0, +\infty]$  be a measure. Let A be an element of  $\mathcal{R}(S)$  with two representations:

$$A = \biguplus_{i=1}^{n} A_i = \biguplus_{j=1}^{p} B_j$$

as a finite union of pairwise disjoint elements of S.

- 1. For i = 1, ..., n, show that  $\mu(A_i) = \sum_{j=1}^p \mu(A_i \cap B_j)$
- 2. Show that  $\sum_{i=1}^n \mu(A_i) = \sum_{j=1}^p \mu(B_j)$
- 3. Explain why we can define a map  $\bar{\mu}: \mathcal{R}(\mathcal{S}) \to [0, +\infty]$  as:

$$\bar{\mu}(A) \stackrel{\triangle}{=} \sum_{i=1}^{n} \mu(A_i)$$

4. Show that  $\bar{\mu}(\emptyset) = 0$ .

EXERCISE 12. Everything being as before, suppose that  $(A_n)_{n\geq 1}$  is a sequence of pairwise disjoint elements of  $\mathcal{R}(\mathcal{S})$ , each  $A_n$  having the representation:

$$A_n = \biguplus_{k=1}^{p_n} A_n^k \ , \ n \ge 1$$

as a finite union of disjoint elements of S. Suppose moreover that  $A = \bigoplus_{n=1}^{+\infty} A_n$  is an element of  $\mathcal{R}(S)$  with representation  $A = \bigoplus_{j=1}^{p} B_j$ , as a finite union of pairwise disjoint elements of S.

- 1. Show that for  $j=1,\ldots,p,$   $B_j=\cup_{n=1}^{+\infty}\cup_{k=1}^{p_n}(A_n^k\cap B_j)$  and explain why  $B_j$  is of the form  $B_j=\uplus_{m=1}^{+\infty}C_m$  for some sequence  $(C_m)_{m\geq 1}$  of pairwise disjoint elements of  $\mathcal{S}$ .
- 2. Show that  $\mu(B_j) = \sum_{n=1}^{+\infty} \sum_{k=1}^{p_n} \mu(A_n^k \cap B_j)$

- 3. Show that for  $n \ge 1$  and  $k = 1, ..., p_n$ ,  $A_n^k = \bigoplus_{j=1}^p (A_n^k \cap B_j)$
- 4. Show that  $\mu(A_n^k) = \sum_{j=1}^p \mu(A_n^k \cap B_j)$
- 5. Recall the definition of  $\bar{\mu}$  of exercise (11) and show that it is a measure on  $\mathcal{R}(\mathcal{S})$ .

EXERCISE 13. Prove the following theorem:

**Theorem 2** Let S be a semi-ring on  $\Omega$ . Let  $\mu: S \to [0, +\infty]$  be a measure on S. There exists a unique measure  $\bar{\mu}: \mathcal{R}(S) \to [0, +\infty]$  such that  $\bar{\mu}_{|S} = \mu$ .

**Definition 10** We define an outer-measure on  $\Omega$  as being any map  $\mu^*$ :  $\mathcal{P}(\Omega) \to [0, +\infty]$  with the following properties:

$$(i) \qquad \mu^*(\emptyset) = 0$$

(ii) 
$$A \subseteq B \Rightarrow \mu^*(A) \le \mu^*(B)$$

(iii) 
$$\mu^* \left( \bigcup_{n=1}^{+\infty} A_n \right) \le \sum_{n=1}^{+\infty} \mu^* (A_n)$$

EXERCISE 14. Show that  $\mu^*(A \cup B) \leq \mu^*(A) + \mu^*(B)$ , where  $\mu^*$  is an outer-measure on  $\Omega$  and  $A, B \subseteq \Omega$ .

**Definition 11** Let  $\mu^*$  be an outer-measure on  $\Omega$ . We define:

$$\Sigma(\mu^*) \stackrel{\triangle}{=} \{ A \subseteq \Omega : \ \mu^*(T) = \mu^*(T \cap A) + \mu^*(T \cap A^c) \ , \ \forall T \subseteq \Omega \}$$

We call  $\Sigma(\mu^*)$  the  $\sigma$ -algebra associated with the outer-measure  $\mu^*$ .

Note that the fact that  $\Sigma(\mu^*)$  is indeed a  $\sigma$ -algebra on  $\Omega$ , remains to be proved. This will be your task in the following exercises.

EXERCISE 15. Let  $\mu^*$  be an outer-measure on  $\Omega$ . Let  $\Sigma = \Sigma(\mu^*)$  be the  $\sigma$ -algebra associated with  $\mu^*$ . Let  $A, B \in \Sigma$  and  $T \subseteq \Omega$ 

- 1. Show that  $\Omega \in \Sigma$  and  $A^c \in \Sigma$ .
- 2. Show that  $\mu^*(T \cap A) = \mu^*(T \cap A \cap B) + \mu^*(T \cap A \cap B^c)$
- 3. Show that  $T \cap A^c = T \cap (A \cap B)^c \cap A^c$
- 4. Show that  $T \cap A \cap B^c = T \cap (A \cap B)^c \cap A$
- 5. Show that  $\mu^*(T \cap A^c) + \mu^*(T \cap A \cap B^c) = \mu^*(T \cap (A \cap B)^c)$
- 6. Adding  $\mu^*(T \cap (A \cap B))$  on both sides 5., conclude that  $A \cap B \in \Sigma$ .
- 7. Show that  $A \cup B$  and  $A \setminus B$  belong to  $\Sigma$ .

EXERCISE 16. Everything being as before, let  $A_n \in \Sigma, n \geq 1$ . Define  $B_1 = A_1$  and  $B_{n+1} = A_{n+1} \setminus (A_1 \cup \ldots \cup A_n)$ . Show that the  $B_n$ 's are pairwise disjoint elements of  $\Sigma$  and that  $\bigcup_{n=1}^{+\infty} A_n = \bigoplus_{n=1}^{+\infty} B_n$ .

EXERCISE 17. Everything being as before, show that if  $B, C \in \Sigma$  and  $B \cap C = \emptyset$ , then  $\mu^*(T \cap (B \uplus C)) = \mu^*(T \cap B) + \mu^*(T \cap C)$  for any  $T \subseteq \Omega$ .

EXERCISE 18.Everything being as before, let  $(B_n)_{n\geq 1}$  be a sequence of pairwise disjoint elements of  $\Sigma$ , and let  $B \stackrel{\triangle}{=} \bigcup_{n=1}^{+\infty} B_n$ . Let  $N \geq 1$ .

- 1. Explain why  $\biguplus_{n=1}^{N} B_n \in \Sigma$
- 2. Show that  $\mu^*(T \cap (\bigcup_{n=1}^N B_n)) = \sum_{n=1}^N \mu^*(T \cap B_n)$
- 3. Show that  $\mu^*(T \cap B^c) \leq \mu^*(T \cap (\biguplus_{n=1}^N B_n)^c)$
- 4. Show that  $\mu^*(T \cap B^c) + \sum_{n=1}^{+\infty} \mu^*(T \cap B_n) \le \mu^*(T)$ , and:
- 5.  $\mu^*(T) \le \mu^*(T \cap B^c) + \mu^*(T \cap B) \le \mu^*(T \cap B^c) + \sum_{n=1}^{+\infty} \mu^*(T \cap B_n)$
- 6. Show that  $B \in \Sigma$  and  $\mu^*(B) = \sum_{n=1}^{+\infty} \mu^*(B_n)$ .
- 7. Show that  $\Sigma$  is a  $\sigma$ -algebra on  $\Omega$ , and  $\mu_{|\Sigma}^*$  is a measure on  $\Sigma$ .

**Theorem 3** Let  $\mu^* : \mathcal{P}(\Omega) \to [0, +\infty]$  be an outer-measure on  $\Omega$ . Then  $\Sigma(\mu^*)$ , the so-called  $\sigma$ -algebra associated with  $\mu^*$ , is indeed a  $\sigma$ -algebra on  $\Omega$  and  $\mu^*_{|\Sigma(\mu^*)}$ , is a measure on  $\Sigma(\mu^*)$ .

EXERCISE 19. Let  $\mathcal{R}$  be a ring on  $\Omega$  and  $\mu : \mathcal{R} \to [0, +\infty]$  be a measure on  $\mathcal{R}$ . For all  $T \subseteq \Omega$ , define:

$$\mu^*(T) \stackrel{\triangle}{=} \inf \left\{ \sum_{n=1}^{+\infty} \mu(A_n) , (A_n) \text{ is an } \mathcal{R}\text{-cover of } T \right\}$$

where an  $\mathcal{R}$ -cover of T is defined as any sequence  $(A_n)_{n\geq 1}$  of elements of  $\mathcal{R}$  such that  $T\subseteq \bigcup_{n=1}^{+\infty}A_n$ . By convention inf  $\emptyset \stackrel{\triangle}{=} +\infty$ .

- 1. Show that  $\mu^*(\emptyset) = 0$ .
- 2. Show that if  $A \subseteq B$  then  $\mu^*(A) \leq \mu^*(B)$ .
- 3. Let  $(A_n)_{n\geq 1}$  be a sequence of subsets of  $\Omega$ , with  $\mu^*(A_n) < +\infty$  for all  $n \geq 1$ . Given  $\epsilon > 0$ , show that for all  $n \geq 1$ , there exists an  $\mathcal{R}$ -cover  $(A_n^p)^{p\geq 1}$  of  $A_n$  such that:

$$\sum_{p=1}^{+\infty} \mu(A_n^p) < \mu^*(A_n) + \epsilon/2^n$$

Why is it important to assume  $\mu^*(A_n) < +\infty$ .

4. Show that there exists an  $\mathcal{R}$ -cover  $(R_k)$  of  $\bigcup_{n=1}^{+\infty} A_n$  such that:

$$\sum_{k=1}^{+\infty} \mu(R_k) = \sum_{n=1}^{+\infty} \sum_{p=1}^{+\infty} \mu(A_n^p)$$

- 5. Show that  $\mu^*(\bigcup_{n=1}^{+\infty} A_n) \le \epsilon + \sum_{n=1}^{+\infty} \mu^*(A_n)$
- 6. Show that  $\mu^*$  is an outer-measure on  $\Omega$ .

EXERCISE 20. Everything being as before, Let  $A \in \mathcal{R}$ . Let  $(A_n)_{n\geq 1}$  be an  $\mathcal{R}$ -cover of A and put  $B_1 = A_1 \cap A$ , and:

$$B_{n+1} \stackrel{\triangle}{=} (A_{n+1} \cap A) \setminus ((A_1 \cap A) \cup \ldots \cup (A_n \cap A))$$

- 1. Show that  $\mu^*(A) \leq \mu(A)$ .
- 2. Show that  $(B_n)_{n\geq 1}$  is a sequence of pairwise disjoint elements of  $\mathcal{R}$  such that  $A= \biguplus_{n=1}^{+\infty} B_n$ .
- 3. Show that  $\mu(A) \leq \mu^*(A)$  and conclude that  $\mu_{|\mathcal{R}}^* = \mu$ .

EXERCISE 21. Everything being as before, Let  $A \in \mathcal{R}$  and  $T \subseteq \Omega$ .

- 1. Show that  $\mu^*(T) \leq \mu^*(T \cap A) + \mu^*(T \cap A^c)$ .
- 2. Let  $(T_n)$  be an  $\mathcal{R}$ -cover of T. Show that  $(T_n \cap A)$  and  $(T_n \cap A^c)$  are  $\mathcal{R}$ -covers of  $T \cap A$  and  $T \cap A^c$  respectively.
- 3. Show that  $\mu^*(T \cap A) + \mu^*(T \cap A^c) \leq \mu^*(T)$ .
- 4. Show that  $\mathcal{R} \subseteq \Sigma(\mu^*)$ .
- 5. Conclude that  $\sigma(\mathcal{R}) \subseteq \Sigma(\mu^*)$ .

EXERCISE 22. Prove the following theorem:

Theorem 4 (Caratheodory's extension) Let  $\mathcal{R}$  be a ring on  $\Omega$  and  $\mu$ :  $\mathcal{R} \to [0, +\infty]$  be a measure on  $\mathcal{R}$ . There exists a measure  $\mu' : \sigma(\mathcal{R}) \to [0, +\infty]$  such that  $\mu'_{\mathcal{R}} = \mu$ .

EXERCISE 23. Let S be a semi-ring on  $\Omega$ . Show that  $\sigma(\mathcal{R}(S)) = \sigma(S)$ .

EXERCISE 24. Prove the following theorem:

**Theorem 5** Let S be a semi-ring on  $\Omega$  and  $\mu: S \to [0, +\infty]$  be a measure on S. There exists a measure  $\mu': \sigma(S) \to [0, +\infty]$  such that  $\mu'_{|S} = \mu$ .

# Solutions to Exercises

#### Exercise 1.

• Let  $x \in A \cap B$ . Then  $x \in B$ . So  $x \notin A \setminus B$ . It follows that  $x \in A \setminus (A \setminus B)$ , and  $A \cap B \subseteq A \setminus (A \setminus B)$ . Let  $x \in A \setminus (A \setminus B)$ . Then  $x \in A$  and  $x \notin A \setminus B$ . But  $x \notin A \setminus B$  implies that either  $x \notin A$  or  $x \in B$ . Hence,  $x \in B$ . finally,  $x \in A \cap B$  and  $A \setminus (A \setminus B) \subseteq A \cap B$ . We have proved that  $A \cap B = A \setminus (A \setminus B)$ 

• Let  $\mathcal{R}$  be a ring and  $A, B \in \mathcal{R}$ . From (iii) of definition (7),  $A \setminus B \in \mathcal{R}$ . Hence,  $A \setminus (A \setminus B) \in \mathcal{R}$ . It follows from the previous point that  $A \cap B \in \mathcal{R}$ . We have proved that a ring is closed under pairwise intersection.

Exercise 1

Exercise 2. Let  $\mathcal{R}$  be ring on  $\Omega$ . Then (i) of definition (6) is immediately satisfied for  $\mathcal{R}$ . From exercise (1), we know that  $\mathcal{R}$  is closed under finite intersection. So (ii) of definition (6) is satisfied for  $\mathcal{R}$ . Let  $A, B \in \mathcal{R}$ . From (iii) of definition (7),  $A \setminus B \in \mathcal{R}$ . Therefore, if we take n = 1 and  $A_1 = A \setminus B \in \mathcal{R}$ , we see that  $A \setminus B = \bigcup_{i=1}^n A_i$  and (iii) of definition (6) is satisfied for  $\mathcal{R}$ . Finally, having checked (i), (ii) and (iii) of definition (6), we conclude that  $\mathcal{R}$  is a semi-ring on  $\Omega$ . Any ring on  $\Omega$  is therefore also a semi-ring on  $\Omega$ .

Exercise 2

#### Exercise 3.

- $\emptyset \in \mathcal{S}_1$  so (i) of definition (6) is satisfied for  $\mathcal{S}_1$ . If  $A, B \in \mathcal{S}_1$ , then  $A \cap B$  is equal to the empty set (remember that  $A_1$ ,  $A_2$  and  $A_3$  are disjoint), unless A (resp. B) is  $\Omega$  itself, or  $A = B \neq \emptyset$ , in which case  $A \cap B$  is equal to B (resp. A). In any case,  $A \cap B \in \mathcal{S}_1$  and condition (ii) of definition (6) is satisfied for  $\mathcal{S}_1$ . If  $A, B \in \mathcal{S}_1$ , since  $\mathcal{S}_1$  has 5 elements,  $A \setminus B$  is one of 25 cases to consider. It is equal to  $\emptyset$ ,  $(\emptyset \setminus \emptyset, \emptyset \setminus A_i, \emptyset \setminus \Omega, A_i \setminus \Omega, A_i \setminus A_i, \Omega \setminus \Omega)$  in 12 of those cases. It is equal to A itself  $(A_i \setminus \emptyset, A_i \setminus A_j, j \neq i, \Omega \setminus \emptyset)$  in 10 of those cases. The last three cases are  $\Omega \setminus A_1 = A_2 \uplus A_3$ ,  $\Omega \setminus A_2 = A_1 \uplus A_3$  and  $\Omega \setminus A_3 = A_1 \uplus A_2$ . Hence, we see that condition (iii) of definition (6) is satisfied for  $\mathcal{S}_1$ . We have proved that  $\mathcal{S}_1$  is indeed a semi-ring on  $\Omega$ .
- If we put  $B_1 = A_1$  and  $B_2 = A_2 \uplus A_3$ , then  $\Omega = B_1 \uplus B_2$  where  $B_1, B_2$  are distinct from  $\emptyset$  and  $\Omega$ . Moreover,  $S_2 = \{\emptyset, B_1, B_2, \Omega\}$ , and proving that  $S_2$  is a semi-ring on  $\Omega$  is identical to the previous point, but is just a little bit easier...
- $S_1 \cap S_2 = \{\emptyset, A_1, \Omega\}$  (remember that all  $A_i$ 's are not empty and pairwise disjoint, so  $A_3 \neq A_2 \uplus A_3$  and  $A_2 \neq A_2 \uplus A_3$ ). Suppose that  $S_1 \cap S_2$  is a semi-ring on  $\Omega$ . Then from (iii) of definition (6), there exists  $n \geq 0$  and  $B_1, B_2, \ldots, B_n$  in  $S_1 \cap S_2$  such that:

$$\Omega \setminus A_1 = B_1 \uplus \ldots \uplus B_n$$

Since  $A_1$  is assumed to be distinct from  $\Omega$ ,  $\Omega \setminus A_1 \neq \emptyset$ . It follows that  $n \geq 1$  and at least one of the  $B_i$ 's is not empty. If  $B_i = \Omega$  then  $\Omega \setminus A_1 = \Omega$  and this would be a contradiction since  $A_1$  is assumed to be not empty. If  $B_i = A_1$  then  $\Omega \setminus A_1 \supseteq A_1$  would also be a contradiction. Hence, the initial assumption of  $S_1 \cap S_2$  being a semi-ring on  $\Omega$  is absurd.  $S_1 \cap S_2$  fails to be a semi-ring on  $\Omega$ . The purpose of this exercise is to show that contrary to Dynkin systems,  $\sigma$ -algebras and rings (as we shall see in the next exercise), taking intersections of semi-rings does not necessarily create another semi-ring. Hence, no attempt will be made to define the notion of generated semi-ring...

Exercise 3

Exercise 4. Each  $\mathcal{R}_i$  being a ring on  $\Omega$ ,  $\emptyset \in \mathcal{R}_i$ . This being true for all  $i \in I$ ,  $\emptyset \in \cap_{i \in I} \mathcal{R}_i = \mathcal{R}$ , and condition (i) of definition (7) is satisfied for  $\mathcal{R}$ . Let  $A, B \in \mathcal{R}$ . Then for all  $i \in I$ , A, B belong to  $\mathcal{R}_i$ . It follows that  $A \setminus B$  and  $A \cup B$  belong to  $\mathcal{R}_i$ . This being true for all  $i \in I$ , both  $A \setminus B$  and  $A \cup B$  lie in  $\cap_{i \in I} \mathcal{R}_i$ , and conditions (ii) and (iii) of definition (7) are satisfied for  $\mathcal{R}$ . Having checked (i), (ii) and (iii) of definition (7), we conclude that  $\mathcal{R}$  is indeed a ring on  $\Omega$ . The purpose of this exercise is to show that an arbitrary (non-empty) intersection of rings on  $\Omega$ , is still a ring on  $\Omega$ .

Exercise 4

# Exercise 5.

- $\emptyset$  being a subset of  $\Omega$ ,  $\emptyset \in \mathcal{P}(\Omega)$  and condition (i) of definition (7) is satisfied for  $\mathcal{P}(\Omega)$ . Given two subsets A, B of  $\Omega$ ,  $A \setminus B$  and  $A \cup B$  are still subsets of  $\Omega$ , i.e.  $A \setminus B \in \mathcal{P}(\Omega)$  and  $A \cup B \in \mathcal{P}(\Omega)$ . Hence, conditions (ii) and (iii) of definition (7) are satisfied for  $\mathcal{P}(\Omega)$ . It follows that  $\mathcal{P}(\Omega)$  is a ring on  $\Omega$ .
- By assumption,  $A \subseteq \mathcal{P}(\Omega)$ . Moreover,  $\mathcal{P}(\Omega)$  is a ring on  $\Omega$ . Therefore,  $\mathcal{P}(\Omega) \in R(A)$ . In particular, R(A) is not empty.
- $\mathcal{R}(\mathcal{A})$  is a non-empty intersection of rings on  $\Omega$ . From exercise (4), it is therefore a ring on  $\Omega$ .
- For all  $\mathcal{R} \in R(\mathcal{A})$ ,  $\mathcal{A} \subseteq \mathcal{R}$ . Hence:

$$\mathcal{A} \subseteq \bigcap_{\mathcal{R} \in R(\mathcal{A})} \mathcal{R} \stackrel{\triangle}{=} \mathcal{R}(\mathcal{A})$$

• Suppose  $\mathcal{R}$  is another ring on  $\Omega$ , with  $\mathcal{A} \subseteq \mathcal{R}$ . Then, by definition of the set  $R(\mathcal{A})$ ,  $\mathcal{R} \in R(\mathcal{A})$ . It follows that:

$$\mathcal{R}(\mathcal{A}) \stackrel{\triangle}{=} \bigcap_{\mathcal{R}' \in R(\mathcal{A})} \mathcal{R}' \subseteq \mathcal{R}$$

So  $\mathcal{R}(\mathcal{A})$  is indeed the *smallest ring* on  $\Omega$  which contains  $\mathcal{A}$ .

Exercise 5

# Exercise 6.

- 1. If  $x \in A_i \cap B_j$  for some i = 1, ..., n and j = 1, ..., p, then  $x \in A \cap B$ . Conversely if  $x \in A \cap B$ , then  $n \ge 1$ ,  $p \ge 1$ , and there exist  $i \in \{1, ..., n\}$  and  $j \in \{1, ..., p\}$  such that  $x \in A_i \cap B_j$ . So  $A \cap B = \bigcup_{i,j} A_i \cap B_j$ . Suppose (i,j) and (i',j') are such that  $(A_i \cap B_j) \cap (A_{i'} \cap B_{j'}) \ne \emptyset$ . In particular,  $A_i \cap A_{i'} \ne \emptyset$ . Since the  $A_i$ 's are pairwise disjoint, we have i = i' and similarly j = j'. Hence, we see that the  $(A_i \cap B_j)_{i,j}$ 's are pairwise disjoint, and finally  $A \cap B = \bigcup_{i,j} A_i \cap B_j$ . From (ii) of definition (6), all the  $A_i \cap B_j$ 's lie in the semi-ring  $\mathcal{S}$ , and we see that  $A \cap B$  is also an element of  $\mathcal{R}$ . We have proved that  $\mathcal{R}$  is closed under finite intersection.
- 2. Since the  $A_i$ 's are pairwise disjoint, for all  $j \in \{1, \ldots, p\}$  being given, the  $A_i \setminus B_j$   $i = 1, \ldots, n$ , are also pairwise disjoint. Hence, the union  $\bigcup_{i=1}^n A_i \setminus B_j$  can legitimately be written as  $\biguplus_{i=1}^n A_i \setminus B_j$ . let  $x \in A \setminus B$ . Then  $x \notin B$ . Thus, for all  $j = 1, \ldots, p$ ,  $x \notin B_j$ . But  $x \in A$ . So there exists  $i \in \{1, \ldots, n\}$  such that  $x \in A_i$ . It follows that for all  $j \in \{1, \ldots, p\}$ ,  $x \in A_i \setminus B_j$  for some  $i \in \{1, \ldots, n\}$ . So  $x \in \bigcap_{j=1}^p \biguplus_{i=1}^n (A_i \setminus B_j)$ . Conversely, suppose that  $x \in \bigcap_{j=1}^p \biguplus_{i=1}^n (A_i \setminus B_j)$ . Then for all  $j \in \{1, \ldots, p\}$ , there exists  $i_j \in \{1, \ldots, n\}$  such that  $x \in A_{i_j} \setminus B_j$ . Since we have assumed  $p \geq 1$ , in particular  $x \in A_{i_1} \subseteq A$ , and for all  $j \in \{1, \ldots, p\}$ ,  $x \notin B_j$ , so  $x \notin B$ . It follows that  $x \in A \setminus B$ . We have proved that:

$$A \setminus B = \bigcap_{i=1}^p \uplus_{i=1}^n (A_i \setminus B_j)$$

- 3. If p = 0, then  $B = \emptyset$  and  $A \setminus B = A \in \mathcal{R}$ . We assume that  $p \geq 1$ . From the previous point, we know that  $A \setminus B = \cap_{j=1}^p C_j$  where  $C_j$  is defined as  $C_j = \bigcup_{i=1}^n A_i \setminus B_j$ . But each  $A_i$  and  $B_j$  is an element of the semi-ring  $\mathcal{S}$ . From (iii) of definition (6), each  $A_i \setminus B_j$  can be written as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . It follows that  $C_j$  itself can be written as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . Hence, we see that for all  $j \in \{1, \ldots, p\}$ ,  $C_j$  is an element of  $\mathcal{R}$ . From 1. we know that  $\mathcal{R}$  is closed under finite intersection. We conclude that  $A \setminus B = \cap_{j=1}^p C_j \in \mathcal{R}$ . We have proved that  $\mathcal{R}$  is closed under pairwise difference.
- 4. Let  $x \in A \cup B$ . then  $x \in A$  or  $x \in B$ . If  $x \in B$  then  $x \in A \setminus B \uplus B$ . If  $x \notin B$  then  $x \in A \setminus B$ . In any case,  $x \in A \setminus B \uplus B$ , and  $A \cup B \subseteq A \setminus B \uplus B$ . Conversely,  $A \setminus B \subseteq A$ , so  $A \setminus B \uplus B \subseteq A \cup B$ . Now, if  $A, B \in \mathcal{R}$ , from the previous point,  $A \setminus B \in \mathcal{R}$ . It follows that  $A \setminus B$  can be written as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . But B itself (being an element of  $\mathcal{R}$ ), can be written as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . It follows that  $A \setminus B \uplus B$  is also a finite union of pairwise disjoint elements of  $\mathcal{S}$ , hence an element of  $\mathcal{R}$ . From  $A \cup B = A \setminus B \uplus B$ , we conclude that  $A \cup B$  is an element of  $\mathcal{R}$ . We have proved that  $\mathcal{R}$  is closed under finite union. Finally, (i), (ii), (iii) of definition (7) being satisfied for  $\mathcal{R}$ ,  $\mathcal{R}$  is indeed a ring on  $\Omega$ .

5. Let  $A \in \mathcal{S}$ . A can obviously be written as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . (Take n=1,  $A_1=A \in \mathcal{S}$  and  $A= \uplus_{i=1}^n A_i$ ). Hence,  $A \in \mathcal{R}$  and  $\mathcal{S} \subseteq \mathcal{R}$ . Consequently, from exercise (5) and the fact that  $\mathcal{R}$  is a ring on  $\Omega$ ,  $\mathcal{R}(\mathcal{S}) \subseteq \mathcal{R}$ . Conversely, let  $A \in \mathcal{R}$ . Then  $A= \uplus_{i=1}^n A_i$  for some  $n \geq 0$  and  $A_i \in \mathcal{S}$ . Since  $\mathcal{S} \subseteq \mathcal{R}(\mathcal{S})$  (see exercise (5)), each  $A_i$  lies in  $\mathcal{R}(\mathcal{S})$ . But from (ii) of definition (7),  $\mathcal{R}(\mathcal{S})$  being a ring is closed under finite union. Hence,  $A \in \mathcal{R}(\mathcal{S})$  and we have  $\mathcal{R} \subseteq \mathcal{R}(\mathcal{S})$ . We have proved that  $\mathcal{R}(\mathcal{S}) = \mathcal{R}$ . The purpose of this exercise is to show that the ring  $\mathcal{R}(\mathcal{S})$  generated by a semi-ring  $\mathcal{S}$  on  $\Omega$ , is equal to the set of all finite unions of pairwise disjoint elements of  $\mathcal{S}$ .

Exercise 6

Exercise 7. Any finite union of pairwise disjoint elements of S, is in particular a finite union of elements of S... So  $R \subseteq R'$ . Let  $A \in R'$ . There exists  $n \geq 0$  and  $A_i \in S$  for i = 1, ..., n such that  $A = \bigcup_{i=1}^n A_i$ . If n = 0, then  $A = \emptyset \in R$ . If  $n \geq 1$ , since  $S \subseteq R = R(S)$ , all  $A_i$ 's are elements of R. R being closed under finite union (it is a ring on  $\Omega$ ), A is itself an element of R. Hence  $R' \subseteq R$ . We have proved that R = R' = R(S). The purpose of this exercise is to show that the generated ring R(S) of a semi-ring S on  $\Omega$ , is also equal to the set of all finite unions of (not necessarily pairwise disjoint) elements of S.

Exercise 7

**Exercise 8.** If  $\mathcal{A}$  is a  $\sigma$ -algebra on  $\Omega$ , then  $A_n \in \mathcal{A}$  and  $A = \bigoplus_{n=1}^{+\infty} A_n$  automatically implies that  $A \in \mathcal{A}$ . Hence, the l.h.s of (ii) and (ii)' are equivalent, whenever  $\mathcal{A}$  is a  $\sigma$ -algebra on  $\Omega$ .

Exercise 8

#### Exercise 9.

1. Define the sequence  $(B_n)_{n\geq 1}$  of elements of  $\mathcal{A}$ , by  $B_i = A_i$  for all  $i = 1, \ldots, n$  and  $B_k = \emptyset$  for all k > n. Then  $A = \bigoplus_{k=1}^{\infty} B_k$ , and since  $A \in \mathcal{A}$ , from (ii) of definition (9), we have:

$$\mu(A) = \sum_{k=1}^{+\infty} \mu(B_k)$$

But from (i) of definition (9),  $\mu(B_k) = 0$  for all k > n. Hence:

$$\mu(A) = \mu(A_1) + \ldots + \mu(A_n)$$

In view of this property, it is customary to say that a measure is *finitely* additive.

2. Suppose  $A, B \in \mathcal{A}$  with  $A \subseteq B$  and  $B \setminus A \in \mathcal{A}$ . Then, we have  $B = A \cup B = A \uplus (B \setminus A)$ . From the previous point we conclude:

$$\mu(A) \le \mu(A) + \mu(B \setminus A) = \mu(B)$$

Exercise 9

Exercise 10.

1. If  $A = \emptyset$ , then either n = 0 or  $A_i = \emptyset$  for all i = 1, ..., n. In any case,  $\bar{\mu}(A) = \sum_{i=1}^n \mu(A_i)$  is true. If  $A \neq \emptyset$ , then  $n \geq 1$ . Since  $\mathcal{S} \subseteq \mathcal{R}(\mathcal{S})$ , all sets involved in  $A = \biguplus_{i=1}^n A_i$  are elements of  $\mathcal{R}(\mathcal{S})$ . Since  $\bar{\mu}$  is a measure on  $\mathcal{R}(\mathcal{S})$ , from exercise (9) we have  $\bar{\mu}(A) = \sum_{i=1}^n \bar{\mu}(A_i)$ . By assumption,  $\bar{\mu}_{|\mathcal{S}} = \mu$  and  $A_i \in \mathcal{S}$  for all i = 1, ..., n. Hence,  $\bar{\mu}(A_i) = \mu(A_i)$  for all i = 1, ..., n. It follows that  $\bar{\mu}(A) = \sum_{i=1}^n \mu(A_i)$ .

2. Let  $A \in \mathcal{R}(\mathcal{S})$ . Then A has a representation  $A = \bigoplus_{i=1}^n A_i$  as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . From the previous point,  $\bar{\mu}(A) = \sum_{i=1}^n \mu(A_i)$ . If  $\bar{\mu}'$  is another measure on  $\mathcal{R}(\mathcal{S})$  with  $\bar{\mu}'_{|\mathcal{S}} = \mu$ , then similarly we have  $\bar{\mu}'(A) = \sum_{i=1}^n \mu(A_i)$ . So  $\bar{\mu}(A) = \bar{\mu}'(A)$ . This being true for all  $A \in \mathcal{R}(\mathcal{S})$ ,  $\bar{\mu} = \bar{\mu}'$ . The purpose of this exercise is to show that if a measure  $\mu$  on a semi-ring  $\mathcal{S}$  can be extended to its generated ring  $\mathcal{R}(\mathcal{S})$ , then such extension is unique.

Exercise 10

# Exercise 11.

1. If p=0, then  $A=\emptyset$ . Then either n=0 and there is nothing to prove, or  $n\geq 1$  with all  $A_i$ 's equal to the empty set. In any case,  $\mu(A_i)=\sum_{j=1}^p \mu(A_i\cap B_j)$  is true. Hence we can assume that  $p\geq 1$ . Since  $A_i\subseteq A$ :

$$A_i = A_i \cap A = \biguplus_{j=1}^p A_i \cap B_j \tag{1}$$

Since S is a semi-ring, it is closed under finite intersection (definition (6)), hence all sets involved in (1) are elements of S. From exercise (9), and the fact that  $\mu$  is a measure on S, we conclude that  $\mu(A_i) = \sum_{j=1}^p \mu(A_i \cap B_j)$ .

2. Similarly to the previous point, for all  $j=1,\ldots,p$  we have  $\mu(B_j)=\sum_{i=1}^n \mu(A_i\cap B_j)$ . It follows that:

$$\sum_{i=1}^{n} \mu(A_i) = \sum_{i=1}^{n} \sum_{j=1}^{p} \mu(A_i \cap B_j) = \sum_{j=1}^{p} \sum_{i=1}^{n} \mu(A_i \cap B_j) = \sum_{j=1}^{p} \mu(B_j)$$

3. Suppose we want to define a map  $\bar{\mu}: \mathcal{R}(\mathcal{S}) \to [0, +\infty]$  with:

$$\bar{\mu}(A) \stackrel{\triangle}{=} \sum_{i=1}^{n} \mu(A_i) \tag{2}$$

where  $A = \bigoplus_{i=1}^n A_i$  is a representation of A as a finite union of pairwise disjoint elements of S. The problem is that such representation may not be unique. However, if  $A = \bigoplus_{j=1}^p B_j$  is another representation of A in terms of finite union of pairwise disjoint elements of S, then from 2.,  $\sum_{i=1}^n \mu(A_i) = \sum_{j=1}^p \mu(B_j)$ . It follows that whichever representation is considered, the sum involved in (2) will still be the same. In other words, definition (2) is unambiguous, and therefore legitimate.

4.  $\emptyset$  has a representation with n = 0, or n = 1 with  $A_1 = \emptyset$ , or n = 2 with  $A_1 = A_2 = \emptyset$ ... Whichever representation we choose for  $\emptyset$ , definition (2) leads to  $\bar{\mu}(\emptyset) = 0$ .

Exercise 11

# Exercise 12.

1. For all j = 1, ..., p, since  $B_j \subseteq A$ , we have:

$$B_j = A \cap B_j = \bigcup_{n=1}^{+\infty} (A_n \cap B_j) = \bigcup_{n=1}^{+\infty} \bigcup_{k=1}^{p_n} (A_n^k \cap B_j)$$

Consider the set  $I = \{(n,k) : n \geq 1, 1 \leq k \leq p_n\}$ . Being a countable union of finite sets, I is a countable set. Hence, there exists a one-to-one map  $\phi : \{m : m \geq 1\} \to I$ . Given  $m \geq 1$ , define  $C_m = A_n^k \cap B_j$  where  $(n,k) = \phi(m)$ . Then we have  $B_j = \bigcup_{m=1}^{+\infty} C_m$ . Since all  $A_n^k$ 's and  $B_j$  itself are elements of the semi-ring S, all  $C_m$ 's are elements of S. Suppose  $C_m \cap C_{m'} \neq \emptyset$  for some  $m,m' \geq 1$ . Then in particular,  $A_n^k \cap A_{n'}^{k'} \neq \emptyset$ , where we have put  $(n,k) = \phi(m)$  and  $(n',k') = \phi(m')$ . Since  $A_n^k \subseteq A_n$  and  $A_{n'}^{k'} \subseteq A_{n'}$ , it follows that  $A_n \cap A_{n'} \neq \emptyset$ , and the  $A_n$ 's being pairwise disjoint, we see that n = n'. Thus,  $A_n^k \cap A_n^{k'} \neq \emptyset$ . But the  $A_n^k$ 's for  $k = 1, \ldots, p_n$  are also pairwise disjoint. We conclude that k = k' and  $\phi(m) = (n,k) = (n',k') = \phi(m')$ . Since  $\phi$  is one-to-one, m = m', and we have proved that  $(C_m)_{m \geq 1}$  is a sequence of pairwise disjoint elements of S.

2. In the previous point, we saw that  $B_j = \bigoplus_{m=1}^{+\infty} C_m$ . Since all sets involved are elements of S and  $\mu$  is a measure on S, from (ii) of definition (9), we have:

$$\mu(B_j) = \sum_{m=1}^{+\infty} \mu(C_m) = \sum_{(n,k)\in I} \mu(A_n^k \cap B_j) = \sum_{n=1}^{+\infty} \sum_{k=1}^{p_n} \mu(A_n^k \cap B_j)$$
 (3)

3. For  $n \ge 1$  and  $k \in \{1, \dots, p_n\}$ , we have  $A_n^k \subseteq A_n \subseteq A$ . Hence:

$$A_n^k = A_n^k \cap A = \biguplus_{j=1}^p (A_n^k \cap B_j)$$

4. From the previous point, using exercise (9), we obtain:

$$\mu(A_n^k) = \sum_{j=1}^p \mu(A_n^k \cap B_j) \tag{4}$$

5. In exercise (11), we saw that the map  $\bar{\mu}: \mathcal{R}(\mathcal{S}) \to [0, +\infty]$  is such that  $\bar{\mu}(\emptyset) = 0$ . Hence (i) of definition (9) is satisfied for  $\bar{\mu}$ . Moreover, by

definition,  $\bar{\mu}(A) = \sum_{j=1}^{p} \mu(B_j)$ . Using equation (3), we have:

$$\bar{\mu}(A) = \sum_{j=1}^{p} \sum_{n=1}^{+\infty} \sum_{k=1}^{p_n} \mu(A_n^k \cap B_j) = \sum_{n=1}^{+\infty} \sum_{k=1}^{p_n} \sum_{j=1}^{p} \mu(A_n^k \cap B_j)$$

Using equation (4), it follows that:

$$\bar{\mu}(A) = \sum_{n=1}^{+\infty} \sum_{k=1}^{p_n} \mu(A_n^k)$$

But, for all  $n \ge 1$ ,  $\bar{\mu}(A_n) = \sum_{k=1}^{p_n} \mu(A_n^k)$ , by definition of  $\bar{\mu}$ . Hence:

$$\bar{\mu}(A) = \sum_{n=1}^{+\infty} \bar{\mu}(A_n)$$

It follows that (ii) of definition (9) is satisfied for  $\bar{\mu}$ . Finally,  $\bar{\mu}$  is a measure on the ring  $\mathcal{R}(\mathcal{S})$ .

Exercise 12

#### Exercise 13.

- Uniqueness is a consequence of exercise (10)
- Take  $\bar{\mu}: \mathcal{R}(\mathcal{S}) \to [0, +\infty]$  as defined in exercise (11). We proved in exercise (12) that  $\bar{\mu}$  is indeed a measure on the ring  $\mathcal{R}(\mathcal{S})$ . Moreover, given  $A \in \mathcal{S}$ , if we take n = 1 and  $A_1 = A$ , then  $A = \bigcup_{i=1}^n A_i$  is a representation of A as a finite union of pairwise disjoint elements of  $\mathcal{S}$ . By definition of  $\bar{\mu}$  (see exercise (11)), it follows that  $\bar{\mu}(A) = \mu(A)$ . This being true for all  $A \in \mathcal{S}$ , we have  $\bar{\mu}_{|\mathcal{S}} = \mu$ . This shows the existence of  $\bar{\mu}$ , and theorem (2) is proved.

Exercise 13

**Exercise 14.** Let  $(A_n)_{n\geq 1}$  be the sequence of subsets of  $\Omega$  defined by  $A_1=A$ ,  $A_2=B$  and  $A_n=\emptyset$  for all  $n\geq 3$ . Using (i) and (iii) of definition (10), we obtain:

$$\mu^*(A \cup B) \le \mu^*(A) + \mu^*(B)$$

Exercise 14

# Exercise 15.

- 1.  $\mu^*$  being an outer measure on  $\Omega$ , by (i) of definition (10), we have  $\mu^*(\emptyset) = 0$ . It follows that given an arbitrary  $T \subseteq \Omega$ ,  $\mu^*(T) = \mu^*(T \cap \Omega) + \mu^*(T \cap \Omega^c)$  is obviously true. Hence, from definition (11),  $\Omega \in \Sigma(\mu^*) = \Sigma$ . The fact that  $A^c \in \Sigma$  is an immediate consequence of definition (11).
- 2. Since  $B \in \Sigma$ , using definition (11) with  $T \cap A$  in place of T, we obtain:

$$\mu^*(T \cap A) = \mu^*(T \cap A \cap B) + \mu^*(T \cap A \cap B^c)$$

3. Since  $A \cap B \subseteq A$ , we have  $A^c \subseteq (A \cap B)^c$ , and consequently:

$$T \cap A^c \subseteq T \cap (A \cap B)^c$$

It follows that:

$$T \cap A^c = (T \cap (A \cap B)^c) \cap T \cap A^c = T \cap (A \cap B)^c \cap A^c$$

4. From  $(A \cap B)^c \cap A = (A^c \cup B^c) \cap A = A \cap B^c$ , we obtain:

$$T \cap (A \cap B)^c \cap A = T \cap A \cap B^c$$

5. Using 3. and 4., we see that the sum  $\mu^*(T \cap A^c) + \mu^*(T \cap A \cap B^c)$  can be expressed as:

$$\mu^*(T \cap (A \cap B)^c \cap A^c) + \mu^*(T \cap (A \cap B)^c \cap A)$$

Since  $A \in \Sigma$ , using definition (11) with  $T \cap (A \cap B)^c$  in place of T, we obtain:

$$\mu^*(T \cap A^c) + \mu^*(T \cap A \cap B^c) = \mu^*(T \cap (A \cap B)^c)$$
 (5)

6. Adding  $\mu^*(T \cap (A \cap B))$  on both sides of equation (5), it appears that the sum:

$$\mu^*(T \cap A^c) + \mu^*(T \cap A \cap B^c) + \mu^*(T \cap A \cap B)$$

is equal to:

$$\mu^*(T \cap (A \cap B)^c) + \mu^*(T \cap (A \cap B))$$

Since  $B \in \Sigma$ , using definition (11) with  $T \cap A$  in place of T, we obtain:

$$\mu^*(T \cap A^c) + \mu^*(T \cap A) = \mu^*(T \cap (A \cap B)^c) + \mu^*(T \cap (A \cap B))$$

and finally, since  $A \in \Sigma$ :

$$\mu^*(T) = \mu^*(T \cap (A \cap B)^c) + \mu^*(T \cap (A \cap B))$$

This being true for all  $T \subseteq \Omega$ , it follows that  $A \cap B \in \Sigma$ . We have proved that  $\Sigma = \Sigma(\mu^*)$  is closed under finite intersection.

7. From  $A \cup B = (A^c \cap B^c)^c$  and the fact that  $\Sigma$  is closed under complementation and finite intersection, we have  $A \cup B \in \Sigma$ . Similarly,  $A \setminus B = A \cap B^c \in \Sigma$ . The purpose of this exercise is to show that the so-called  $\sigma$ -algebra  $\Sigma(\mu^*)$  associated with an outer measure  $\mu^*$ , is closed under finite intersection and union, and closed under complementation and difference.

Exercise 15

#### Exercise 16.

• Suppose  $n \geq 1$ ,  $p \geq 1$  and  $B_n \cap B_p \neq \emptyset$ . Without loss of generality, we can assume that  $n \leq p$ . Suppose n < p and  $x \in B_n \cap B_p$ . Since  $x \in B_n$ , we have  $x \in A_n$ . However, since  $x \in B_p$ ,  $x \notin A_1 \cup \ldots \cup A_{p-1}$ . In particular,  $x \notin A_n$ . This is a contradiction. It follows that if  $B_n \cap B_p \neq \emptyset$  then n = p, and  $(B_n)_{n \geq 1}$  is a sequence of pairwise disjoint subsets of  $\Omega$ .

- From exercise (15), all  $B_n$ 's are in fact elements of  $\Sigma$ .
- Since for all  $n \geq 1$ ,  $B_n \subseteq A_n$ , we have:  $\bigoplus_{n=1}^{+\infty} B_n \subseteq \bigcup_{n=1}^{+\infty} A_n$ . Conversely, suppose  $x \in \bigcup_{n=1}^{+\infty} A_n$ . Then, there exists  $n \geq 1$  such that  $x \in A_n$ . Consider the set:

$$I(x) \stackrel{\triangle}{=} \{ n \ge 1, x \in A_n \}$$

This set is a non-empty subset of  $\mathbf{N}^*$  (the set of all positive integers). It follows that I(x) has a smallest element p. If p=1, then  $x\in A_1=B_1$ . If p>1, then  $x\in A_p\setminus (A_1\cup\ldots\cup A_{p-1})=B_p$ . In any case,  $x\in B_p\subseteq \uplus_{n=1}^{+\infty}B_n$ . Consequently, it follows that  $\cup_{n=1}^{+\infty}A_n\subseteq \uplus_{n=1}^{+\infty}B_n$ .

• We have proved that  $(B_n)_{n\geq 1}$  is a sequence of pairwise disjoint elements of  $\Sigma$ , such that:

$$\bigcup_{n=1}^{+\infty} A_n = \biguplus_{n=1}^{+\infty} B_n$$

Exercise 16

**Exercise 17.** Let  $B, C \in \Sigma$  be such that  $B \cap C = \emptyset$ . Since  $B \in \Sigma$ , using definition (11) with  $T \cap (B \uplus C)$  in place of T, we have:

$$\mu^*(T\cap (B\uplus C))=\mu^*(T\cap (B\uplus C)\cap B)+\mu^*(T\cap (B\uplus C)\cap B^c)$$

From  $B \cap C = \emptyset$  and in particular  $C \subseteq B^c$ , we obtain:

$$\mu^*(T \cap (B \uplus C)) = \mu^*(T \cap B) + \mu^*(T \cap C)$$

Note that it was not necessary to use the fact that both B and C were elements of  $\Sigma$ .

Exercise 17

# Exercise 18.

- 1.  $\biguplus_{n=1}^{N} B_n \in \Sigma$  is an immediate consequence of exercise (15).
- 2. Using exercise (17) with a simple induction argument, we obtain:

$$\mu^*(T \cap (\uplus_{n=1}^N B_n)) = \sum_{n=1}^N \mu^*(T \cap B_n)$$

3. Since  $\bigcup_{n=1}^{N} B_n \subseteq B$ , we have  $T \cap B^c \subseteq T \cap (\bigcup_{n=1}^{N} B_n)^c$ . Using (ii) of definition (10), we obtain:

$$\mu^*(T \cap B^c) \le \mu^*(T \cap (\biguplus_{n=1}^N B_n)^c)$$

4. Using 2. and 3., if we put  $C_N = \bigcup_{n=1}^N B_n$ , we have:

$$\mu^*(T \cap B^c) + \sum_{n=1}^N \mu^*(T \cap B_n) \le \mu^*(T \cap (C_N)^c) + \mu^*(T \cap C_N)$$

However from 1.,  $C_N \in \Sigma$ . Using definition (11), we obtain:

$$\mu^*(T \cap B^c) + \sum_{n=1}^N \mu^*(T \cap B_n) \le \mu^*(T)$$

Taking the limit as  $N \to +\infty$ , we conclude:

$$\mu^*(T \cap B^c) + \sum_{n=1}^{+\infty} \mu^*(T \cap B_n) \le \mu^*(T)$$

5. Since  $T = (T \cap B^c) \cup (T \cap B)$ , using exercise (14):

$$\mu^*(T) \le \mu^*(T \cap B^c) + \mu^*(T \cap B)$$

However,  $T \cap B = \bigcup_{n=1}^{+\infty} T \cap B_n$ . Using (iii) of definition (10), we have:

$$\mu^*(T \cap B) \le \sum_{n=1}^{+\infty} \mu^*(T \cap B_n)$$

It follows that:

$$\mu^*(T) \le \mu^*(T \cap B^c) + \mu^*(T \cap B) \le \mu^*(T \cap B^c) + \sum_{n=1}^{+\infty} \mu^*(T \cap B_n)$$

6. From 4. and 5., we see that  $\mu^*(T) = \mu^*(T \cap B^c) + \mu^*(T \cap B)$ . This being true for all  $T \subseteq \Omega$ , it follows that  $B = \biguplus_{n=1}^{+\infty} B_n \in \Sigma$ . Also, from 4. and 5., we have:

$$\mu^*(T) = \mu^*(T \cap B^c) + \sum_{n=1}^{+\infty} \mu^*(T \cap B_n)$$

In particular, taking T = B, using the fact that  $\mu^*(\emptyset) = 0$ , we obtain:

$$\mu^*(B) = \sum_{n=1}^{+\infty} \mu^*(B_n)$$

7. We saw in exercise (15) that  $\Sigma$  contains  $\Omega$ , and is closed under complementation. If  $(A_n)_{n\geq 1}$  is a sequence of elements of  $\Sigma$ , then from exercise (16), there exists a sequence  $(B_n)_{n\geq 1}$  of pairwise disjoint elements of  $\Sigma$ , with  $B= \biguplus_{n=1}^{+\infty} B_n= \bigcup_{n=1}^{+\infty} A_n$ . In 6., we saw that such B is an element of  $\Sigma$ . It follows that  $\bigcup_{n=1}^{+\infty} A_n \in \Sigma$ , and  $\Sigma$  is closed under countable union. Hence, we have proved that  $\Sigma$  is a  $\sigma$ -algebra on  $\Omega$ .  $\mu^*$  being an outer measure on  $\Omega$ ,  $\mu^*(\emptyset)=0$ . So (i) of definition (9) is satisfied for  $\mu^*_{|\Sigma}$ . If  $(B_n)_{n\geq 1}$  is a sequence of pairwise disjoint elements of  $\Sigma$ , and  $B=\biguplus_{n=1}^{+\infty} B_n$ , we saw in 6. that:

$$\mu^*(B) = \sum_{n=1}^{+\infty} \mu^*(B_n)$$

It follows that (ii) of definition (9) is satisfied for  $\mu_{|\Sigma}^*$ . Finally,  $\mu_{|\Sigma}^*$  is indeed a measure on  $\Sigma$ . The purpose of the exercise is to prove theorem (3).

Exercise 18

# Exercise 19.

1.  $\mathcal{R}$  being a ring on  $\Omega$ ,  $\emptyset \in \mathcal{R}$ . If we define a sequence  $(A_n)_{n\geq 1}$ , with  $A_n = \emptyset$  for all  $n \geq 1$ , then  $(A_n)_{n\geq 1}$  is an  $\mathcal{R}$ -cover of the empty set. It follows that:

$$\mu^*(\emptyset) \le \sum_{n=1}^{+\infty} \mu(A_n) = 0$$

Moreover,  $\mu^*(\emptyset)$  being the infimum over a set of non-negative numbers, we have  $\mu^*(\emptyset) \geq 0$ . Finally  $\mu^*(\emptyset) = 0$ .

2. Let  $A \subseteq B \subseteq \Omega$ . Let  $(B_n)_{n\geq 1}$  be an  $\mathcal{R}$ -cover of B. Then in particular,  $(B_n)_{n\geq 1}$  is an  $\mathcal{R}$ -cover of A. It follows that:

$$\mu^*(A) \le \sum_{n=1}^{+\infty} \mu(B_n) \tag{6}$$

Hence,  $\mu^*(A)$  is a lower bound of all sums involved in (6), as  $(B_n)_{n\geq 1}$  ranges over all  $\mathcal{R}$ -covers of B.  $\mu^*(B)$  being the infimum of those sums, it is the greatest of such lower bounds, from which we conclude that  $\mu^*(A) \leq \mu^*(B)$ .

3. Since  $\mu^*(A_n) < +\infty$ , we have  $\mu^*(A_n) < \mu^*(A_n) + \epsilon/2^n$ . It follows that  $\mu^*(A_n) + \epsilon/2^n$  cannot be a lower bound of all sums  $\sum_{p=1}^{+\infty} \mu(B_p)$ , as  $(B_p)_{p\geq 1}$  ranges over all  $\mathcal{R}$ -covers of  $A_n$ . Hence, there exists an  $\mathcal{R}$ -cover  $(A_n^p)^{p\geq 1}$  of  $A_n$  such that:

$$\sum_{p=1}^{+\infty} \mu(A_n^p) < \mu^*(A_n) + \frac{\epsilon}{2^n}$$

It is important to assume  $\mu^*(A_n) < +\infty$ , since otherwise the inequality  $\mu^*(A_n) \leq \mu^*(A_n) + \epsilon/2^n$  may not be a strict inequality, and the above reasoning would fail.

4.  $\mathbf{N}^*$  being the set of positive integers,  $\mathbf{N}^* \times \mathbf{N}^*$  is a countable set. There exists a one-to-one map  $\phi: \mathbf{N}^* \to \mathbf{N}^* \times \mathbf{N}^*$ . Given  $k \geq 1$ , define  $R_k = A_n^p$ , where  $(n,p) = \phi(k)$ . Then  $(R_k)_{k \geq 1}$  is a sequence of elements of  $\mathcal{R}$  such that:

$$\bigcup_{n=1}^{+\infty} A_n \subseteq \bigcup_{n=1}^{+\infty} \bigcup_{p=1}^{+\infty} A_n^p = \bigcup_{k=1}^{+\infty} R_k$$

In other words,  $(R_k)_{k\geq 1}$  is an  $\mathcal{R}$ -cover of  $\bigcup_{n=1}^{+\infty} A_n$ . Moreover:

$$\sum_{k=1}^{+\infty} \mu(R_k) = \sum_{(n,p) \in \mathbf{N}^* \times \mathbf{N}^*} \mu(A_n^p) = \sum_{n=1}^{+\infty} \sum_{p=1}^{+\infty} \mu(A_n^p)$$

5. It follows from 4. that:

$$\mu^*(\bigcup_{n=1}^{+\infty} A_n) \le \sum_{k=1}^{+\infty} \mu(R_k) = \sum_{n=1}^{+\infty} \sum_{p=1}^{+\infty} \mu(A_n^p)$$

Hence, using 3.:

$$\mu^*(\bigcup_{n=1}^{+\infty} A_n) \le \sum_{n=1}^{+\infty} (\mu^*(A_n) + \frac{\epsilon}{2^n})$$

and finally:

$$\mu^*(\cup_{n=1}^{+\infty} A_n) \le \epsilon + \sum_{n=1}^{+\infty} \mu^*(A_n)$$
 (7)

6. From 1. and 2., we see that (i) and (ii) of definition (10) are satisfied for  $\mu^*$ . Let  $(A_n)_{n\geq 1}$  be a sequence of subsets of  $\Omega$ . If  $\mu^*(A_n)=+\infty$  for some  $n\geq 1$ , then:

$$\mu^*(\cup_{n=1}^{+\infty} A_n) \le \sum_{n=1}^{+\infty} \mu^*(A_n)$$
 (8)

is obviously true. If  $\mu^*(A_n) < +\infty$  for all  $n \geq 1$ , then given  $\epsilon > 0$  from 5., inequality (7) holds. Since  $\epsilon$  is arbitrary, it follows that inequality (8) still holds. Hence, (iii) of definition (10) is satisfied for  $\mu^*$ . Finally,  $\mu^*$  is an outer-measure on  $\Omega$ .

Exercise 19

18

# Exercise 20.

1. Since  $A \in \mathcal{R}$ , the sequence  $(R_n)_{n\geq 1}$  defined by  $R_1 = A$  and  $R_n = \emptyset$  for all  $n \geq 2$ , is an  $\mathcal{R}$ -cover of A. Hence:

$$\mu^*(A) \le \sum_{n=1}^{+\infty} \mu(R_n) = \mu(A)$$

2. Suppose  $n \geq 1$ ,  $p \geq 1$  and  $B_n \cap B_p \neq \emptyset$ . Without loss of generality, we can assume that  $n \leq p$ . Suppose n < p and  $x \in B_n \cap B_p$ . Since  $x \in B_n$ , we have  $x \in A_n \cap A$ . However, since  $x \in B_p$ ,  $x \notin (A_1 \cap A) \cup \ldots \cup (A_{p-1} \cap A)$ . In particular,  $x \notin A_n \cap A$ . This is a contradiction. It follows that if  $B_n \cap B_p \neq \emptyset$  then n = p, and  $(B_n)_{n \geq 1}$  is a sequence of pairwise disjoint subsets of  $\Omega$ . From exercise (1), we know that a ring is closed under finite intersection. From (ii) and (iii) of definition (7), it is also closed under finite union and difference. It follows that all  $B_n$ 's are in fact elements of  $\mathcal{R}$ . Since for all  $n \geq 1$ ,  $B_n \subseteq A_n \cap A$ , we have:

$$\biguplus_{n=1}^{+\infty} B_n \subseteq \bigcup_{n=1}^{+\infty} A_n \cap A = A \cap \bigcup_{n=1}^{+\infty} A_n = A$$

Conversely, suppose  $x \in A \subseteq \bigcup_{n=1}^{+\infty} A_n$ . Then, there exists  $n \ge 1$  such that  $x \in A_n \cap A$ . Consider the set:

$$I(x) \stackrel{\triangle}{=} \{ n \ge 1, x \in A_n \cap A \}$$

This set is a non-empty subset of  $\mathbf{N}^*$  (the set of all positive integers). It follows that I(x) has a smallest element p. If p=1, then  $x\in A_1\cap A=B_1$ . If p>1, then by definition of p, we have  $x\in (A_p\cap A)\setminus ((A_1\cap A)\cup\ldots\cup (A_{p-1}\cap A))=B_p$ . In any case,  $x\in B_p\subseteq \bigoplus_{n=1}^{+\infty}B_n$ . Consequently, it follows that  $A\subseteq \bigoplus_{n=1}^{+\infty}B_n$ . We have proved that  $(B_n)_{n\geq 1}$  is a sequence of pairwise disjoint elements of  $\mathcal{R}$ , such that:  $A=\bigoplus_{n=1}^{+\infty}B_n$ 

3.  $\mu$  being a measure on  $\mathcal{R}$ , from 2. we obtain:

$$\mu(A) = \sum_{n=1}^{+\infty} \mu(B_n)$$

Since for all  $n \geq 1$ , we have  $B_n \subseteq A_n$ , it follows from exercise (9) that  $\mu(B_n) \leq \mu(A_n)$ . Hence:

$$\mu(A) \le \sum_{n=1}^{+\infty} \mu(A_n) \tag{9}$$

The  $\mathcal{R}$ -cover  $(A_n)_{n\geq 1}$  of A being arbitrary, we see that  $\mu(A)$  is a lower bound of all sums involved in (9), as  $(A_n)_{n\geq 1}$  ranges across all  $\mathcal{R}$ -covers of A.  $\mu^*(A)$  being the greatest of such lower bounds, it follows that  $\mu(A) \leq \mu^*(A)$ . Using 1., we conclude that  $\mu(A) = \mu^*(A)$ . This being true for all  $A \in \mathcal{R}$ , we have proved that  $\mu^*_{|\mathcal{R}} = \mu$ .

Exercise 20

#### Exercise 21.

1. We saw in exercise (19) that  $\mu^*$  is an outer measure on  $\Omega$ . From exercise (14), and the fact that  $T = (T \cap A) \cup (T \cap A^c)$ , we obtain:

$$\mu^*(T) \le \mu^*(T \cap A) + \mu^*(T \cap A^c)$$

2. If  $(T_n)_{n\geq 1}$  is an  $\mathcal{R}$ -cover of T, then in particular  $T_n \in \mathcal{R}$  for all  $n\geq 1$ . Since  $A\in \mathcal{R}$ , it follows from exercise (1) that  $T_n\cap A\in \mathcal{R}$ , and from (iii) of definition (7) that  $T_n\cap A^c=T_n\setminus A\in \mathcal{R}$ , for all  $n\geq 1$ . Moreover, from  $T\subseteq \cup_{n=1}^{+\infty} T_n$ , we have:

$$T \cap A \subseteq \bigcup_{n=1}^{+\infty} T_n \cap A$$
$$T \cap A^c \subseteq \bigcup_{n=1}^{+\infty} T_n \cap A^c$$

We conclude that  $(T_n \cap A)_{n\geq 1}$  and  $(T_n \cap A^c)_{n\geq 1}$  are  $\mathcal{R}$ -covers of  $T \cap A$  and  $T \cap A^c$  respectively.

3. It follows from 2. that:

$$\mu^*(T \cap A) \le \sum_{n=1}^{+\infty} \mu(T_n \cap A)$$

$$\mu^*(T \cap A^c) \le \sum_{n=1}^{+\infty} \mu(T_n \cap A^c)$$

However,  $\mu$  being a measure on  $\mathcal{R}$ , from exercise (9), we have:

$$\mu(T_n) = \mu(T_n \cap A) + \mu(T_n \cap A^c)$$

for all  $n \geq 1$ . It follows that:

$$\mu^*(T \cap A) + \mu^*(T \cap A^c) \le \sum_{n=1}^{+\infty} \mu(T_n)$$

This being true for all  $\mathcal{R}$ -covers  $(T_n)_{n\geq 1}$  of T, we finally have:

$$\mu^*(T \cap A) + \mu^*(T \cap A^c) \le \mu^*(T)$$

4. Given  $A \in \mathcal{R}$ , we see from 1. and 3. that for all  $T \subseteq \Omega$ :

$$\mu^*(T) = \mu^*(T \cap A) + \mu^*(T \cap A^c)$$

Hence, from definition (11), it follows that A is an element of  $\Sigma(\mu^*)$ , (the  $\sigma$ -algebra associated with the outer measure  $\mu^*$ ). This being true for all  $A \in \mathcal{R}$ , we have proved that  $\mathcal{R} \subseteq \Sigma(\mu^*)$ .

5. The  $\sigma$ -algebra  $\sigma(\mathcal{R})$  generated by  $\mathcal{R}$ , is the smallest  $\sigma$ -algebra on  $\Omega$  containing  $\mathcal{R}$ . Thus, it follows immediately from 4. that  $\sigma(\mathcal{R}) \subseteq \Sigma(\mu^*)$ .

Exercise 21

# Exercise 22.

- Let  $\mu': \sigma(\mathcal{R}) \to [0, +\infty]$  be defined by  $\mu' = \mu_{|\sigma(\mathcal{R})}^*$ , where  $\mu^*$  is the outer measure on  $\Omega$  defined in exercise (19). We saw in exercise (20) that  $\mu_{|\mathcal{R}}^* = \mu$ . Hence, since  $\mathcal{R} \subseteq \sigma(\mathcal{R})$ , we have  $\mu_{|\mathcal{R}}' = \mu_{|\mathcal{R}}^* = \mu$ .
- From theorem (3), we know that  $\mu^*_{|\Sigma(\mu^*)}$  is a measure on  $\Sigma(\mu^*)$ . However,  $\sigma(\mathcal{R}) \subseteq \Sigma(\mu^*)$  (exercise (21)). It is an immediate consequence of definition (9), that if we restrict the measure  $\mu^*_{|\Sigma(\mu^*)}$  to the smaller  $\sigma$ -algebra  $\sigma(\mathcal{R})$ , the resulting map is a measure defined on  $\sigma(\mathcal{R})$ . But the restriction of  $\mu^*_{|\Sigma(\mu^*)}$  to  $\sigma(\mathcal{R})$  is nothing but  $\mu'$ . It follows that  $\mu'$  is indeed a measure on  $\sigma(\mathcal{R})$ . This proves theorem (4).

Exercise 22

**Exercise 23.** Let S be a semi-ring on  $\Omega$ . Since  $S \subseteq \mathcal{R}(S) \subseteq \sigma(\mathcal{R}(S))$ , we have  $\sigma(S) \subseteq \sigma(\mathcal{R}(S))$ . However,  $S \subseteq \sigma(S)$ . Moreover, from exercise (7),  $\mathcal{R}(S)$  is the set of all finite unions of elements of S. Since the  $\sigma$ -algebra  $\sigma(S)$ 

is in particular closed under finite union, it follows that  $\mathcal{R}(\mathcal{S}) \subseteq \sigma(\mathcal{S})$  and consequently  $\sigma(\mathcal{R}(\mathcal{S})) \subseteq \sigma(\mathcal{S})$ . Finally, we have proved that  $\sigma(\mathcal{R}(\mathcal{S})) = \sigma(\mathcal{S})$ . Exercise 23

**Exercise 24.** From theorem (2), the measure  $\mu: \mathcal{S} \to [0, +\infty]$  can be extended to the ring  $\mathcal{R}(\mathcal{S})$  generated by  $\mathcal{S}$ . In other words, there exists a measure  $\bar{\mu}: \mathcal{R}(\mathcal{S}) \to [0, +\infty]$  such that  $\bar{\mu}_{|\mathcal{S}} = \mu$ . From theorem (4), the measure  $\bar{\mu}: \mathcal{R}(\mathcal{S}) \to [0, +\infty]$  can be extended the  $\sigma$ -algebra  $\sigma(\mathcal{R}(\mathcal{S}))$  generated by  $\mathcal{R}(\mathcal{S})$ . In other words, there exists a measure  $\mu': \sigma(\mathcal{R}(\mathcal{S})) \to [0, +\infty]$ , such that  $\mu'_{|\mathcal{R}(\mathcal{S})|} = \bar{\mu}$ . However, from exercise (23),  $\sigma(\mathcal{R}(\mathcal{S})) = \sigma(\mathcal{S})$ . Moreover, since  $\mathcal{S} \subseteq \mathcal{R}(\mathcal{S})$ , we have  $\mu'_{|\mathcal{S}} = \bar{\mu}_{|\mathcal{S}} = \mu$ . It follows that  $\mu'$  is a measure on  $\sigma(\mathcal{S})$  such that  $\mu'_{|\mathcal{S}} = \mu$ . This proves theorem (5).

Exercise 24